

Integral Consulting Inc. 285 Century Place Suite 190 Louisville, CO 80027

telephone: 303.404.2944 facsimile: 303.404.2945 www.integral-corp.com

MEMORANDUM

To: Jason Gunter, U.S. Environmental Protection Agency

From: Marcia Greenblatt

Date: February 2014

Subject: Big River OU-2 Sediment Probing Survey

Project No.: C816

Sediment probing in Big River, Owl Creek, and Flat River Creek was performed from November 29 to December 1, 2012 to support the ongoing feasibility study in operable unit 2 (OU-2) (river and non-residential areas) of St. Francois County Mining Area (also referred to as Big River Mine Tailings Site), Missouri. The sediment probing survey was performed to support the identification and evaluation of potential remedial alternatives. The primary objective of sediment probing was to estimate current sediment volumes in depositional areas and in tributaries. The proposed approach to sediment probing was documented in the Field Sampling Proposal (FSP), Appendix A of the Quality Assurance Project Plan (QAPP) (Integral 2013a).

SUMMARY OF SEDIMENT PROBING ACTIVITIES

A sediment probing survey was performed to estimate sediment volumes in depositional areas and tributaries within the former mining areas. Targeted locations included 1) pooled or backwater areas to characterize sediment trapping potential and 2) tributaries previously surveyed in 2007 to estimate current sediment volumes (NewFields 2007). Locations for sediment probing were identified following a review of the Visual River Survey (VRS) observations and previous sediment volume surveys (Integral 2013b).

A review of the VRS observations indicated only one significant Big River backwater area where sediments may collect within OU-2, which was located at the Bone Hole. Three transects were performed at the Bone Hole: BRFSSP01, BRFSSP02, and BRFSSP03 (Figure 1, Table 1). Mining-related material was previously observed in two tributaries, Flat River Creek and Owl Creek (NewFields 2007), and these tributaries were resurveyed (transects

40460432 Superfund 4.2

Sediment Probing Survey
St. Francois County Mining Area (Big River Mine Tailings Site)
February 2014
Page 2 of 7

FRCSP01 through FRCSP10 and transects OCFSSP01 through OCFSSP03). Two additional Big River sediment probing locations were added to the survey following the submittal of the FSP at the request of the U.S. Environmental Protection Agency (Gunter 2012, pers. comm.): one at St. Francois State Park and one at Blackwell. The Blackwell location was not surveyed because of access restrictions and a water depth that was too deep to safely wade (Integral 2013c). Four sediment probing transects were performed near St. Francois State Park (transects SFSP01 through SFSP04); these transects were performed in areas where depositional material was observed at the time of survey.

Field Approach

To estimate the average thickness of the sediment in the selected depositional areas, 20 cross-sections of Big River and selected tributaries were probed with a 3/8-in., 4.5-ft long rebar probe where sediments were present (Integral 2013a). The number and spacing of the cross-sections were determined based on the extent of the depositional area. At each river cross-section, five to eleven depth estimates were made by pushing the probe by hand from the surface of the sediment to refusal and measuring the penetration depth (Table 2). For the purposes of this study, sediments are defined as all material in the channel that occurs above the refusal depth or what is taken to be intact or broken bedrock (Integral 2013a). This definition includes main channel and off-channel deposits, both in and above the water level. The exposed gravel and sand bars that are common in Big River were included in the sediment probing transects.

The channel width, or cross-section length, was obtained using a hip chain. The length of the string pulled from the hip chain was measured (to the nearest foot) and recorded in the field logbook. Photographs were taken by the field team to document general observations and the condition of each probing area (Integral 2013c; Attachment A). At each location, a form was completed to document the probing transects, including layout, width, depth, and substrate composition information (Attachment B).

Data Evaluation

The probing data were used to estimate sediment volumes for each of the surveyed areas. The unit sediment volume (the volume per unit length) was calculated at each transect as the product of the average probing depth and the transect width (Table 3). The sediment volume between two successive transects (the segment volume) was calculated as the product of the average unit sediment volume of the upstream and downstream transects and segment length (the distance between adjacent transects) (Table 4). Segment lengths were measured along the channel centerline within a Geographic Information System (GIS)

Sediment Probing Survey
St. Francois County Mining Area (Big River Mine Tailings Site)
February 2014
Page 3 of 7

using 2010 National Agriculture Imagery Program (NAIP) aerial photography (NAIP 2010). The sediment volumes for each segment were summed to estimate the total sediment volume in each area surveyed.

Results

The calculated sediment depths and volumes of each study area, as well as observations made during the survey, are briefly discussed below. Observed trends and calculated values are compared with previous sediment probing studies (NewFields 2007, MDNR 2007), and the Borrow Pit Monitoring Project (Owen et al. 2012).

Big River at the Bone Hole

Three sediment probing transects were performed from the confluence of Big River with Owl Creek to the low-water crossing at the Bone Hole (BRFSSP01 through BRFSSP03). Results from transects BRFSSP01 through BRFSSP03 indicated a depositional environment, with average sediment depths along the three transects of 1.9, 1.6, and 1.6 ft respectively (Table 3). The total sediment volume between the transects was estimated to be 547 yd³; the observed deposition is consistent with the low-water crossing slowing the flow of Big River and creating a depositional backwater area.

Collocated transects from the 2007 and 2012 surveys, BR-3 and BRFSSP03, had average sediment depths of a similar magnitude (1.75 and 1.6 ft, respectively), suggesting that the Bone Hole has stable sediment depths over time (Table 3) (NewFields 2007). This finding is consistent with results from the 2010 borrow pit study performed at the Bone Hole. The study was performed to characterize infilling potential following excavation of 508 yd³ of sediment (Owen et al. 2012). Post-excavation surveys indicated that the pit had filled to 80 percent of its previous volume following a 10-year flood event and had nearly completely refilled after several near-bankfull events; it was concluded that the channel sediment storage at the Bone Hole is relatively stable (Owen et al. 2012).

Big River at St. Francois State Park

Four sediment probing transects were performed in Big River adjacent to St. Francois State Park in which the 1,900 ft-long surveyed area contained approximately 22,000 yd³ of sediment (Figure 1, Table 4). Gravel and sand bars were observed in each St. Francois State Park probing transect and contained approximately 70 percent of the estimated volume. Each transect also had locations where sediment was probed to depths greater than 1 ft in the river channel. The sand and gravel bars as well as the river bottom of this stretch of Big River are an area where sediments are stored along Big River.

Sediment Probing Survey St. François County Mining Area (Big River Mine Tailings Site) February 2014 Page 4 of 7

Average sediment depths along the transects adjacent to St. Francois State Park were similar or greater than those observed at the Bone Hole (Table 3). It is likely that the average sediment depths for transects SFSP01 and SFSP02 were underestimated because the probe did not reach refusal for several of the locations along these transects and, therefore, a sediment depth of 60 in. was assumed in the volume calculations (Table 2). Missouri Department of Natural Resources measured sediment depths with ground-penetrating radar (GPR) in a bar approximately 150 ft upstream of transect SFSP01; GPR data indicated sediment depths up to 9.7 ft (MDNR 2007).

Owl Creek

Sediment probing was performed along the lower 1,000 ft of Owl Creek at three transects, OCFSSP01 through OCFSSP03. Little sediment deposition was observed in the uppermost transect, OCFSSP01, and the creek bottom was scoured to bedrock at some locations along this transect (Table 2). Average sediment depth along Transect OCFSSP02 was greater than OCFSSP01, and the substrate primarily consisted of sandy silt and fines. A vegetated bar was observed in the center of OCFSSP02 and leaf litter was observed in Owl Creek at this location. Transect OCFSSP03 had an average sediment depth of 1.5 ft, the deepest of the Owl Creek transects; no sand or gravel bars were observed along OCFSSP03. The 2007 survey occupied approximately the same transects in Owl Creek and found a similar pattern of increasing sediment depths with distance downstream (NewFields 2007).

In 2007, the total estimated sediment volume in Owl Creek was 3,650 yd3 from the culvert at the abandoned Mississippi River and Bonne Terre railroad crossing to the confluence with Big River, approximately 1,200 ft downstream (Figure 1, Table 4) (NewFields 2007). In the current sediment survey this value was 1,206 yd3, indicating approximately a 3-fold decrease in sediment volume. NewFields (2007) used a different approach for volume calculation than in the current study: the Owl Creek transects were used as approximate midpoints to calculate volumes for three segments within this span (NewFields 2007). Applying this approach to the 2012 survey data, the sediment volume from the railroad crossing to the confluence of Owl Creek and Big River was approximately 1,660 yd³, less than one-half of the volume estimated in 2007. In 2007 the water was too deep to collect probing measurements at the lowermost transect (Owl Creek C-C'), and an average sediment depth of 3 ft was assumed (NewFields 2007), which could have led to an overestimate of sediment volume. In 2012 all locations along this transect were successfully probed and the average sediment depth was 1.5, one-half of the 2007 assumed value. Although the sediment volume in Owl Creek may have been overestimated in 2007 because of the estimated depth along the lower transect, the average sediment depths for the two upper Owl Creek transects were also higher in 2007 than in 2012, when the average depths

Sediment Probing Survey
St. Francois County Mining Area (Big River Mine Tailings Site)
February 2014
Page 5 of 7

were less than 1 foot. Sediments observed in the Owl Creek reach in 2007 may have washed off from the small watershed, moved downstream from the upper reach, and potentially eroded from the Mississippi River and Bonne Terre Railroad crossing (NewFields 2007), located approximately 100 ft above the survey reach. Sediments observed in the upper reach of Owl Creek in 2007 were vegetated and appeared stable (NewFields 2007). The reduction in sediment volume could indicate that there is little or no ongoing erosion of the railroad crossing and that sediments have been washed out of the creek.

Flat River Creek

Ten sediment probing transects were performed in Flat River Creek from the drainage of the Elvins/Rivermines pile to the confluence with Big River. The majority of sediment in Flat River Creek was found in gravel and sand bars observed in transects FRCFSSP03, -05, -06 and -10 (Figure 1). Several of these bars were vegetated and appeared stable. Limited sediment was observed in the Flat River Creek channel and the creek bottom was scoured to bedrock in portions of several transects (FRCFSSP01, -02, -04, -05, -07, and -08) (Figure 1, Table 2). The patterns observed in the 2012 sediment probing survey were generally consistent with those observed in the 2007 study, where the only significant sediment accumulation was within gravel and sand bars (NewFields 2007).

The estimated sediment volume in the lower 5 miles of Flat River Creek was 21,000 yd³ in 2012 (Table 4). In 2007, only one complete sediment probing transect was performed (at FR-8 [FRCFSSP10]); measurements of gravel and sand bars observed between FR-4 and FR-7.5 (FRCFSSP01 and FRCFSSP09) were used to estimate a total sediment volume of 13,000 yd³. The observed difference in sediment volume could be due to the different survey techniques.

Source control measures via non-time critical removal actions (NTCRAs) have been established at the Elvins/Rivermines piles; stabilization of the National and Federal piles is ongoing (USEPA 2012). The stabilization measures will prevent mining-related material from entering Big River and its tributaries; completion of the NTCRAs is expected to lead to a decrease in Flat River Creek sediment loading in the future.

Summary

Sediment probing was performed to support sediment volume estimates in several reaches in Big River and its tributaries. The accumulation of sediment within Big River at the locations evaluated suggests that these locations act as long-term sediment traps. The

Sediment Probing Survey
St. François County Mining Area (Big River Mine Tailings Site)
February 2014
Page 6 of 7

effectiveness of trapping and periodically removing sediments as a component of remediation will be evaluated in the Feasibility Study.

Sediment deposition patterns within Owl Creek and Flat River Creek were generally similar between the 2007 and 2012 probing surveys. Sediment was primarily stored in gravel and sand bars, some of which were vegetated and appeared stable. Given the dynamic hydrologic nature of Big River watershed (where flows can rise over an order of magnitude in less than a day), sediment volumes can be expected to vary over time. The ongoing remediation of the piles adjacent to Flat River Creek may result in a reduction of sediment supply following construction completion. Over time, a decline in sediment volume may be observable. Although the sediment load from the Flat River Creek watershed has not been evaluated, the sediment budget will considered as part of the Feasibility Study.

REFERENCES

Gunter, J. 2012. Personal communication (email to M. Nations, Doe Run, and M. Greenblatt, Integral Consulting on October 2, 2012, regarding ??). U.S. Environmental Protection Agency, Washingon DC.

Integral. 2013a. Field sampling plan, river and nonresidential areas (OU-2) St. Francois County Mining Area, Missouri. Appendix A to the quality assurance project plan, river and non-residential areas (OU-2) St. Francois County Mining Area, Missouri. Prepared for The Doe Run Resources Corporation, St. Louis, MO. Integral Consulting Inc., Seattle, WA.

Integral. 2013b. Visual river survey technical memorandum, river and non-residential areas (OU-2) St. Francois County Mining Area, Missouri. Prepared for The Doe Run Resources Corporation, St. Louis, MO. Integral Consulting Inc., Seattle, WA.

Integral. 2013c. Field sampling report, river and non-residential areas (OU-2) St. Francois County mining area, Missouri. Prepared for The Doe Run Resources Corporation, St. Louis, MO. Integral Consulting Inc., Seattle, WA.

MDNR. 2007. The estimated volume of mine-related benthic sediment in Big River at two point bars in St. Francois State Park using ground penetrating radar and x-ray fluorescence. Missouri Department of Natural Resources, Field Services Division, Water Quality Monitoring Section, Jefferson City, MO.

NAIP. 2010. Digital orthorectified images (DOQ) for St. Francois County, Missouri. Available at: http://msdis.missouri.edu/data/naip2010/index.html. Accessed October 4,

Sediment Probing Survey
St. Francois County Mining Area (Big River Mine Tailings Site)
February 2014
Page 7 of 7

2011. U.S. Department of Agriculture, Farm Service Agency, Aerial Photography Field Office, National Agricultural Imagery Program (NAIP), Salt Lake City, UT.

NewFields. 2007. Volume of sediments in Big River, Flat River Creek, and Owl Creek. Prepared for The Doe Run Resources Corporation, St. Louis, MO. NewFields, Denver, CO.

Owen, M.R., Pavlowsky R.T., and D.J. Martin. 2012. Big River Borrow Pit monitoring project. Big River Mining Sediment Project. Funded by U.S. Fish and Wildlife Service and Cooperative Ecosystems Studies Unit. Prepared by the Ozarks Environmental and Water Resources Institute, Missouri State University, Springfield, MO.

USEPA. 2012. NPL site description for Big River mine tailings/St. Joe Minerals Corporation Site, Missouri. EPA ID# MOD981126899. Available at: http://www.epa.gov/region07/cleanup/npl_files/mod981126899.pdf. Accessed April 8, 2013. Dated April 24, 2012. U.S. Environmental Protection Agency Region 7, Kansas City, KS.

LIST OF ATTACHMENTS

Figure 1. Sediment Probing Locations

Table 1. Locations and Coordinates of Sediment Probing Survey Transects

Table 2. Sediment Probing Survey Field Measurements


Table 3. Transect Average Sediment Depth and Unit Volume

Table 4. Estimated Sediment Volumes

Attachment A. Photos from 2012 Sediment Probing Survey

Attachment B. Field forms from 2012 Sediment Probing Survey

FIGURES



TABLES

Table 1. Locations and Coordinates of Sediment Probing Survey Transects

| Transect Number | Location | Longitude ^a | Latitude ^a |
|-----------------|--------------------------------------|------------------------|-----------------------|
| BRFSSP01 | Big River at Bone Hole | -90.550700000 | 37.875220000 |
| BRFSSP02 | Big River at Bone Hole | -90.550582000 | 37.875264000 |
| BRFSSP03 | Big River at Bone Hole | -90.550462000 | 37.875297000 |
| SFSP01 | Big River at St. Francois State Park | -90.540230385 | 37.954617066 |
| SFSP02 | Big River at St. Francois State Park | -90.539963788 | 37.954455530 |
| SFSP03 | Big River at St. Francois State Park | -90.539655009 | 37.954313601 |
| SFSP04 | Big River at St. Francois State Park | -90.541207860 | 37.958735462 |
| OCFSSP01 | Owl Creek | -90.552234744 | 37.872289176 |
| OCFSSP02 | Owl Creek | -90.551908703 | 37.873450854 |
| OCFSSP03 | Owl Creek | -90.551181918 | 37.874653350 |
| FRCFSSP01 | Flat River Creek | -90.530685432 | 37.836008944 |
| FRCFSSP02 | Flat River Creek | -90.526226734 | 37.839986267 |
| FRCFSSP03 | Flat River Creek | -90.523338299 | 37.842441910 |
| FRCFSSP04 | Flat River Creek | -90.519660498 | 37.845405664 |
| FRCFSSP05 | Flat River Creek | -90.517131248 | 37.849211154 |
| FRCFSSP06 | Flat River Creek | -90.512417566 | 37.855413540 |
| FRCFSSP07 | Flat River Creek | -90.504091823 | 37.862988454 |
| FRCFSSP08 | Flat River Creek | -90.505106687 | 37.867138518 |
| FRCFSSP09 | Flat River Creek | -90.503051600 | 37.882675244 |
| FRCFSSP10 | Flat River Creek | -90.498998711 | 37.887894044 |

Note

^a Coordinates are from the center of the probing transect and are reported in the World Geodetic System 1984 (WGS 1984).

Table 2. Sediment Probing Survey Field Measurements

| Transect | Bank Width | | | | | Probed Se | diment Dep | oth (inches) | (| | | |
|-----------|------------|---------|---------|-----------------|-----------------|-----------------|------------|-----------------|---------|---------|----------|-----------------------|
| Number | (feet) | Depth 1 | Depth 2 | Depth 3 | Depth 4 | Depth 5 | Depth 6 | Depth 7 | Depth 8 | Depth 9 | Depth 10 | Depth 11 ⁶ |
| BRFSSP01 | 149 | 17 | 4 | 5 | 10 | 23 | 20 | 38 | 36 | 40 | 40 | - |
| BRFSSP02 | 152 | 30 | 31 | 32 | 26 | 6 | 14 | 26 | 15 | 10 | 3 | |
| BRFSSP03 | 173 | 29 | 16 | 22 | 29 | 14 | 19 | 17 | 18 | 22 | 5 | |
| SFSP01 | 113 | 57.5 | 44 | 60 ^b | 60 ^b | 60 ^b | 47 | 60 ^b | 27.5 | 29 | 0 | |
| SFSP02 | 108.5 | 17.5 | 9.5 | 60 ^b | 4.5 | 48 | 46 | 43 | 12 | 0 | 0 | |
| SFSP03 | 120.5 | 24 | 8.5 | 4 | 6 | 23 | 41 | 16 | 11 | 29 | 38 | |
| SFSP04 | 142.5 | 15 | 37 | 3 | 1.5 | 11 | 14 | 31 | 4 | 4 | 49 | |
| OCFSSP01 | 40 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 2 | 0.75 | 1.5 | 5 |
| OCFSSP02 | 38 | 20 | 14.5 | 2.5 | 14.5 | 11.5 | 5 | 3 | 7 | 9.25 | 17 | - |
| OCFSSP03 | 46 | 49 | 18.5 | 17 | 9 | 10.5 | 11 | 6.5 | 4 | 4 | 8 | 59 |
| FRCFSSP01 | 74 | 9 | 7 | 7 | 6 | 0 | 0 | 0 | 0 | 0 | 10.5 | |
| FRCFSSP02 | 59 | 23 | 3.5 | 4.25 | 2 | 4.5 | 4.25 | 0 | 0 | 1.5 | 5 | - |
| FRCFSSP03 | 72 | 2 | 3 | 4 | 2.5 | 4 | 2.75 | 3 | 2 | 5 | 5 | · · |
| FRCFSSP04 | 79.5 | 0.25 | 14.5 | 0 | 0 | 0 | | | | | | |
| FRCFSSP05 | 70 | 7 | 26 | 9.5 | 18 | 11.5 | 29.5 | 3 | 0.5 | 0 | 9.5 | |
| FRCFSSP06 | 73 | 1 | 1.5 | 1.25 | 2 | 1.25 | 3 | 3 | 2 | 3 | 5 | - |
| FRCFSSP07 | 29.5 | 6 | 4 | 3 | 5 | 4.5 | 0 | 0 | 0 | 0 | 0 | - |
| FRCFSSP08 | 40 | 0 | 0 | 0 | 0 | 2 | 5 | 5 | 6.5 | 4.5 | 4 | |
| FRCFSSP09 | 36 | 5.25 | 5.5 | 5.5 | 8 | 3.5 | 4 | 2.5 | 8 | 10 | 14 | |
| FRCFSSP10 | 72 | 4.5 | 5 | 12 | 17.5 | 18 | 6 | 25 | 17 | 16 | 11 | 8.5 |

Notes:

^a A goal of 5 to 10 measurements at a sediment probing transect was proposed in the FSP; at some locations an additional transect was performed for better coverage of the area (Integral 2013c).

b In cases where the probe did not reach refusal, a note was made and a depth value of 60 inches (maximum probe depth) was indicated.

Table 3. Transect Average Sediment Depth and Unit Volume

| Transect Number | Bank Width (ft) | Average Sediment Depth (ft) | Unit Sediment Volume ^a (yd ³ /ft) |
|--------------------|--------------------|--------------------------------|--|
| BRFSSP01 | 149 | 1.9 | 11 |
| BRFSSP02 | 152 | 1.6 | 9.1 |
| BRFSSP03 | 173 | 1.6 | 10 |
| SFSP01 | 113 | 3.7 | 16 |
| SFSP02 | 108.5 | 2.0 | 8.1 |
| SFSP03 | 120.5 | 1.7 | 7.5 |
| SFSP04 | 142.5 | 1.4 | 7.5 |
| OCFSSP01 | 40 | 0.1 | 0.10 |
| OCFSSP02 | 38 | 0.9 | 1.2 |
| OCFSSP03 | 46 | 1.5 | 2.5 |
| FRCFSSP01 | 74 | 0.3 | 0.90 |
| FRCFSSP02 | 59 | 0.4 | 0.87 |
| FRCFSSP03 | 72 | 0.3 | 0.74 |
| FRCFSSP04 | 79.5 | 0.2 | 0.72 |
| FRCFSSP05 | 70 | 1.0 | 2.5 |
| FRCFSSP06 | 73 | 0.2 | 0.52 |
| FRCFSSP07 | 29.5 | 0.2 | 0.20 |
| FRCFSSP08 | 40 | 0.2 | 0.33 |
| FRCFSSP09 | 36 | 0.6 | 0.74 |
| FRCFSSP10 | 72 | 1.1 | 2.8 |

Note:

^a Unit volume is calculated as the sediment volume per unit length.

Table 4. Estimated Sediment Volumes

| Upstream Transect | Downstream Transect | Location | Segment Length ^a (ft) | Volume (yd ³) | 2007 Volume (yd ³) |
|-------------------|------------------------|---|----------------------------------|------------------------------|-----------------------------------|
| BRFSSP01 | BRFSSP02 | Big River at Bone Hole | 30 | 297 | |
| BRFSSP02 | BRFSSP03 | Big River at Bone Hole | 26 | 250 | |
| | | Total for Big River at Bone Hole: | 56 | 547 | |
| SFSP01 | SFSP04 | Big River at St. Francois State Park | 1,697 | 20,558 | |
| SFSP02 | SFSP01 | Big River at St. Francois State Park | 97 | 1,136 | |
| SFSP03 | SFSP02 | Big River at St. Francois State Park | - 103 | 803 | |
| | | Total for Big River at St. Francois State Park: | 1,897 | 22,497 | |
| OCFSSP01 | OCFSSP02 | Owl Creek | 440 | 298 | |
| OCFSSP02 | OCFSSP03 | Owl Creek | 495 | 908 | |
| | | Total for lower Owl Creek: | 935 | 1,206 | 3,650 ^b |
| FRCFSSP01 | FRCFSSP02 | Flat River Creek | 1,986 | 1,783 | |
| FRCFSSP02 | FRCFSSP03 | Flat River Creek | 1,256 | 1,032 | |
| FRCFSSP03 | FRCFSSP04 | Flat River Creek | 1,564 | 1,147 | |
| FRCFSSP04 | FRCFSSP05 | Flat River Creek | 1,664 | 2,764 | |
| FRCFSSP05 | FRCFSSP06 | Flat River Creek | 2,868 | 4,351 | |
| FRCFSSP06 | FRCFSSP07 | Flat River Creek | 3,721 | 1,339 | |
| FRCFSSP07 | FRCFSSP08 | Flat River Creek | 1,622 | 431 | |
| FRCFSSP08 | FRCFSSP09 | Flat River Creek | 7,354 | 4,021 | |
| FRCFSSP09 | FRCFSSP10 | Flat River Creek | 2,439 | 3,943 | |
| | | Total for lower Flat River Creek: | 24,474 | 20,811 | 13,385° |

Notes:

^a Segment length was measured along river or creek centerline in ArcGIS using NAIP aerial photos (NAIP 2010).

^b The 2007 survey defined lower Owl Creek as the reach from the MR&BT railroad crossing to the Big River confluence (NewFields 2007). Surveyed stations were used as approximate midpoints for finding the volume of three segments in this reach. Applying this approach to the 2012 survey data, the sediment volume from the railroad crossing to the confluence of Owl Creek and Big River was approximately 1,660 yd³.

^c The 2007 survey defined lower Flat River Creek as the reach from river mile 5 to the Big River confluence (NewFields 2007). Gravel and sand bars were measured to make sediment volume estimates from station FR-4 (2012 station FRCFSSP01) to FR-7.5 (2012 station FRCFSSP09). The unit volume calculated from station FR-8 measurements (2012 station FRCFSSP10) were used to calculate the volume from station FR-7.5 to the Big River confluence.

ATTACHMENT A

PHOTOS FROM 2012 SEDIMENT PROBING SURVEY

(FOUND ON ACCOMPANYING DISC)



Photo 1: Big River at Bone Hole transect BRFSSP01, right bank in foreground



Photo 2: Big River at Bone Hole transect BRFSSP02, facing the left bank



Photo 3: Big River at Bone Hole transect BRFSSP03, facing the left bank (low water crossing on right)



Photo 4: Big River at Bone Hole transect BRFSSP03, sediment



Photo 5: Big River at St. Francois State Park transect SFSP01, facing upstream



Photo 6: Big River at St. Francois State Park transect SFSP02, facing downstream



Photo 7: Big River at St. Francois State Park transect SFSP03, facing downstream



Photo 8: Big River at St. Francois State Park transect SFSP04, facing downstream



Photo 9: Owl Creek transect OCFSSP01, facing upstream and culvert of abandoned railroad bridge



Photo 10: Owl Creek transect OCFSSP01, facing downstream



Photo 11: Owl Creek transect OCFSSP02, facing downstream




Photo 12: Owl Creek transect OCFSSP03, facing left bank



Photo 13: Flat River Creek transect FRCFSSP01, facing downstream



Photo 14: Flat River Creek transect FRCFSSP02, facing downstream



Photo 15: Flat River Creek transect FRCFSSP03, facing downstream



Photo 16: Flat River Creek transect FRCFSSP04, facing downstream and the right bank



Photo 17: Flat River Creek transect FRCFSSP05, facing downstream



Photo 18: Flat River Creek transect FRCFSSP06, facing left bank, gravel bar in foreground



Photo 19: Flat River Creek transect FRCFSSP07, facing downstream



Photo 20: Flat River Creek transect FRCFSSP08, facing downstream




Photo 21: Flat River Creek transect FRCFSSP09, facing downstream



Photo 22: Flat River Creek transect FRCFSSP10, facing the left bank

ATTACHMENT **B**

FIELD FORMS FROM 2012 SEDIMENT PROBING SURVEY

(FOUND ON ACCOMPANYING DISC)



| | Sample Location: BRFSSPD1 | Date: 1/29/12 Samplers: \$0,132,55 |
|--------------|--|--|
| | Longitude: 10ft bank 37,8704898 N | Date: 1/29 12 Samplers: SD, BL, JJ Right Beink: 90, 55025564 h Latitude: Right bank: 90,55066541 W |
| | Bank Width (m) 149 ft | Number of Sediment Depth Estimates: 10 |
| SD". | Sediment Depths: (5 to 10 measurements) | 2 11 = 211 22 11 11=11 (12+ |
| WD: | 17" 4" 5" 10" 23" - | 20' 38" 36" 40" 40† 38" 35" 37" 40" 41" |
| | Site Sketch | ef Kenterno? wooder |
| 51)=(ediment | | 1 - 10 |
| ub = worter | | |
| depth | | water an road |
| | BIG RIVER | CVOSSING |
| | Flow | |
| | | |
| | 57 | |
| | 7./ | |
| | 100000 | |
| | | gravel Roa |
| | Cross Section (provide both horizontal and ve | ertical measurements) |
| auteur - | Gross Section (provide Both Herizonian and re | January Medical Cinemas, |
| Celler. | 10 9 8 7 6 5 4 | 1321 |
| Wood | ed to the total | Mooded |
| | | |
| | 12 | gravel |
| | | Today |
| | | |
| | Notes (Photo Number and substrate descripti | on): |
| 7 | 35- white board | |
| 3 | 36- Sediment probing | |
| 3 | 8 -sediment probe a 10th state | in an transect |
| 0 |) Solnery ground, 6) Sandy | g more |
| (2) | | Smally Sandy growel |
| 3 | (E) 1/ | Smally Sandy gravel |
| 9 | (1) | |
| 0 | (1) (1) SOFT SITI | y clay |
| | The state of the s | PH 15 (11) |


| Sample Location: BRFSC02 Left Bank: 37:87544724*N Longitude: Lyni Bank: 37:87506476 | Date: 1/29/12 Samplers: SW1BL, S WH Bank: 90.550494751W N Latitude: Right bonk: 90.550491411W |
|--|---|
| Longitude: Fynt Bank: 37. 87506476 | 'N Latitude: Right honk: 90.55049141'W |
| Bank Width (m) 152th | Number of Sediment Depth Estimates:_/O |
| Sediment Depths: (5 to 10 measurements) SD = 30" 31" 32" 26" (0" Site Sketch SD = Sediment depth SD = Sediment de | 144 26" 15" 10" 3" 43" 43" 42" 40" 31" Toolewfoo deep for access from O-15" |
| No=waterdepth BlaRIVER | 15' -0 30' FD low water water cussing over 105' -0 105' -0 105' -0 105' -0 105' -0 |
| Croe v | gravelroad Flore |
| Cross Section (provide both horizontal and | 91 10 June est |
| | |
| | |
| Notes (Photo Number and substrate descri 31-White board 32-looking US a) strand width mo 33-looky DS a) sedim proby | 245 |
| @ sandy graves (8) sand | G grand G grand Stave |

| | Sample Location: BRFSSP03 Left Bank: 37.87557467 N Longitude: Right Bank: 37.87506187 N Bank Width (m) 173.0 Ft | Date: 1/29/17 Samplers: SW 131, 75 Left Bank: 90. 55033414 W Latitude: Right bank: 90. 15060313 |
|--------------------------------|--|--|
| UP! | Sediment Depths: (5 to 10 measurements) 16" 22" 29" 14" 31" 35" 37" Site Sketch | 19" 17" 18" 22" 5" 34" 36" 34" 33" 34" |
| depth 0 = nexter- depth. | BIG RIVER | 173.00 (30 1.00) 3 (20 1.00) 3 |
| | | gravel rond and Flow |
| | Cross Section (provide both horizontal and v | 8 9 10 19 mudline |
| | Notes (Photo Number and substrate descript | refusal tion): |
| | 24-Uniteboard 25-Proping location 03 26-Measuring distance to transect 1 27-Measury wo/sp 38-measing wo/sp | |
| (<u>)</u> | D-sandy graves in | Sandy gravel Sandy gravel Sandy gravel Sandy gravel Sandy gravel Sandy gravel Cobbb / gravel |

| | Sample Location: OCFSSP0 / | Date: 1/29/12 Samplers: 340.500 1) |
|-------|--|--|
| | Longitude: 37.87229187 N | Latitude: 90.55773827-60 |
| | Bank Width (m) 151/101/10 HO FT | Number of Sediment Depth Estimates://_ |
| | Sediment Depths: (58to 10 measurements) | 20" 24 28 32 36 40 |
| 50 = | 0.0 0.0 | 0.0" 0.0" 2.0" 0.75" 1.5" 5.0 |
| w1) > | 5.0" 6.5" 7.0" 10.75" 17.2)" Site Sketch Crossection | 12" 13" 125" 10" 4.25" 5.0" |
| | | |
| | | |
| | | A |
| | p 2 3 4 | 5 6 7 8 9 10 11 |
| | | veg e text |
| | rock | |
| | 1 polps | - Sediment |
| | | depty |
| | | |
| | | |
| | | |
| | Gross Section (provide both horizontal and | vertical measurements) |
| | DIE SLETZE. Wooded + | vegetated my |
| | very etated | 10 CSP 112- |
| | 180 | Flow on Us tanned |
| | | |
| | hauta | by OWLCREEK Stendy flow, |
| | transect | from 12.0 |
| | Roc | kledges. |
| | Notes (Photo Number and substrate descrip | otion): |
| | light algae, al light silt lay | f transect |
| | | |
| | 48-100Kg (B to RB ab) | frahect so - whiteboad |
| ſ. | | ble, Silty sand |
| 15 |) 11 (7) 11 | |
| | | 1 |
| (3 | 10 51/A | sand, coloble |

Sediment Probing Section Form

| | Sample Location: OCFSSP&Z Longitude: 37.87345561 N | Date: $\frac{11/29/12}{29/12}$ Samplers: $\frac{5W}{BU}$ BL $\frac{5}{1}$ Latitude: $\frac{90.55/9/180}{1}$ |
|-----------|--|--|
| | Bank Width (m) 38,6 ft | Number of Sediment Depth Estimates: |
| SiD = | Sediment Depths: (5 to 10 measurements) | 21 24 28 34 37.5 |
| WD- | $\frac{20''}{20''} \frac{145''}{30''} \frac{2.5''}{2.3''} \frac{14.5''}{7.0''} \frac{11.5''}{0.5''}$ | 511 3.0" 7.0" 9.251 17" 0" 7.5" 5.0" 5.75" 3" |
| | Site Sketch 1 wooded Lst | The state of the s |
| 50 = sedu | meet lepth | 4 年 4 |
| (1) = L | at depth | Big River |
| | " 2 | Flore |
| | 9: 0 | |
| | gravet A manthum | t A P |
| | bay wooded ske | ez bark |



Notes (Photo Number and substrate description):

" (Photo Number a

| | Sample Location: OFFSSPØ3 | Date: 11/29/12 Samplers: 500 66 55 |
|------------|--|---|
| | Longitude: 37.87465750 °N | Latitude: 99.551183751W |
| | Bank Width (m) | Number of Sediment Depth Estimates: |
| SD= WD= | Sediment Depths: (5 to 10 measurements) | 1\0" 6.5" 4.0" 4.70" 8.0" 59.0" 33.5"8 280 350 3.5" |
| | The state of the siles | wooded area 1/2 Big |
| | ONEO | 200 |
| | 4 | |
| | 1200 ant | 1000 |
| | 167 | Plan |
| | 3100 | |
| | Te made | Owl Creek |
| | | |
| | 有民 有民公 | auded area AR |
| | | |
| l l | Cross Section (provide both horizontal and v | ertical measurements) |
| | left bank | 5 |
| | | |
| | 00000 | |
| | | 3 |
| | Notes (Photo Number and substrate descript | ion): |
| . (| | ddl of transcat |
| | 40-Jane Simol Standing on right | ounk. |
| 9 | Can be contained to the first | |
| | 0-1-1-0-11-0 | and and Bandon of the |
| | Oclay, hard, soft at the top. (9) | Sandy agravel Occupie when soft sith loose sandy around on top of sandy grave |
| 1 | | |
| 2 | 3 clay | sandy gravel |
| | Blight samu arayol a | and a discher |
| | 3 clay | |





| | ample Location: FRCFSSPØ1 Date: 1/30/12 Samplers: SW, BL, JS | |
|------|--|-----|
| | ongitude: 77,83601039 N Latitude: 90.53068939 N | |
| | ank Width (m) 74ft Number of Sediment Depth Estimates: | |
| | Number of Sediment Depth Estimates: Number of Sediment Depth Estimates: 39' 16 53 74' 74' 74' 74' 74' 74' 74' 74' 75' 75' 75' 75' 75' 75' 75' 75' 75' 75 | |
| 50° | 6'' $9.0''$ $9.0''$ $9.0''$ $9.0''$ $9.0''$ $9.0''$ | |
| cup: | ediment Depths: (5 to 10 measurements) 0' 7.0" 4.0" 6.0" 6.0" 6.0" 6.75" 7.0" 6.75" 11.0" 10.0" 10.0" 10. | |
| | RIPRADSLOPE | |
| | N/IMPC | |
| | SUND FLAT RIVER CREFIL | |
| | | |
| | 2 | |
| | FLAT RIVER CREEK | |
| | CLEEK | |
| | The yegetated flatarea | |
| | | |
| | RIPKAP SLOPE WOODED | |
| | | |
| | * Drawing Snot to scale | |
| | ross Section (provide both horizontal and vertical measurements) | |
| | Like 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 | |
| | moding bedroom | - / |
| | Fluchies Huckness | _ |
| | otes (Photo Number and substrate description): | |
| | 5-White bon 1 88 100kg US from under Bridge | 2 |
| | 6-100 hg D5 | - |
| | 2 - 10/1 0 1 to 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 | |
| | 7-10km 2 transact RB to LB | |
| | grass w/ suit/sand sitty loan v/ Designanics | |

| | Sample Location: FRC F-SP # 2 | Date: 30/12 Samplers: SW, BL, JS |
|-----|---|---|
| | Or william | Latitude: 90.52623824°W |
| | Bank Width (m) 57.011 | Number of Sediment Depth Estimates: 10 |
| | Sediment Depths: (5 to 10 measurements) $= 23" - 3.5" - 4.25" - 2" - 4.5" - 4.5"$ | 130" 22" 23" 10" 10" |
| WV= | 100000 | ded lank |
| | | |
| | FIAT RIVER CREEK | We made made |
| | | |
| | | WOODED (to ~ 60' NW) their by mater trees |
| | -5 | * Drawys not to scale |
| | Cross Section (provide both horizontal and ver | |
| | CB 0 0 0 0 0 0 0 | D 8 9 1 |
| | | |
| | sediment sediment | Sed most |
| | Notes (Photo Number and substrate description Coord collected a culture | |
| | 99 - White board 92. | - 100 My DS. |
| | D Sundy sitt/organio 5)19 | Cherellubble over Sandy SIF |
| | a gravel/Lobble 65 | grandfubble over sandy silt |
| | 3 | sienel sand |
| | (4) | grevel Saver |

| 50= WP= | Sample Location: FRCFSPD3 Longitude: 37.8424469 N Bank Width (m) 72.0 H Sediment Depths: (5 to 10 measurements) 3.0" 4.0" 2.5" 4.0" 2.5" 4.0" 2.5" 5.0" 5.0" 5.0" Site Sketch |) |
|------------|---|------------------|
| | Flat River Freek Greek Gravel Wooded -x x x x x x x x x x x x x x x x x x x | state - fence |
| (b) | Cross Section (provide both horizontal and vertical measurements) Skep Skep What We do not We | |
| | Notes (Photo Number and substrate description): Coprols collected from center of gravel har 93 - White bound 94 - RB to LB also transect 97 - 100 hg DS D raised gravel 95 - 100 hg US Organics (Sconbul/Grave sand (D) 11 (D) 12 (D) 12 (D) 13 (D) 13 (D) 14 (D) 15 (D) 16 (D) 16 | el ba |


| | Sample Location: FRCFSSPØ4 | Date: 1/30/12 Samplers: SAN, BL, JS |
|------------|--|--|
| | Longitude: 37.84540711°N | Latitude: 90.51966549°60 |
| | Bank Width (m) 79.5 H | Number of Sediment Depth Estimates: 5 |
| | Sediment Denths: (5 to 10 measurements) | |
| 50= | 0.15" 14.5" 0.0' 0.0' 0.0' 0.0" 0.0" 7.5" 4" High | |
| LD= | 0.0" 0.0" 7.5" 4" Hig | h St. |
| | Site Sketch High School | |
| | Shallow bank, vege tat | T d |
| SD=SEdi | pts 150 de | |
| 1 0 = 1100 | 1 | redward (15'-60') from co |
| ho=ug | pth bedrock b | from co |
| | SUM C | FLATRIVER |
| | - Gan | CREEK |
| | · | Plow |
| | | to toppes a love and last below raily |
| | Steep | bank, traces a lowend, ballast below fruit |
| | XIXIXIX | line |
| | V- N | + + - / |
| | | awiys not to scale |
| | Cross Section (provide both horizontal and v | ertical measurements) vail for tracks |
| | ing" | 3 D 5 |
| | 13, 0 0 | 3 |
| | | |
| | BEDROCK | water |
| | DEPOLCE | bedrock |
| | Sed mont bal | bedrock |
| | thickness bedrock | |
| | Notes (Photo Number and substrate descript | ion): 8 1/30/12 |
| | Coords Collected 2 centers | Hansect left bank (target location |
| | 70 -11 | |
| | 73 - Whiteboard | 81-100kg US oftranked Deed with |
| | 80 - 11 alox R. | 84-100h DS from bedrock pool 84-10chg RB to LB alvoss transact |
| | Define Cilt | 84- locke RB to LB alross |
| | | |
| | @ Silt w/ veg ste | frank c |
| ı | 3 bedwar | frankl ct |
| į | @ Silt w/ veg ste | frankl ct |





Sediment Probing Section Form

| Sample Location: FRCFSSPØ7 Longitude: 37.86300212 N Bank Width (m) 29.5' Sediment Depths: (5 to 10 measurements) SD = 6.0" 4.5" OO 5.15" 4.5" | Date: $\frac{1}{30}$ rz Samplers: \underline{SW} , \underline{BL} , \underline{TS} Latitude: $\underline{90.50910189}$ W Number of Sediment Depth Estimates: $\underline{10}$ $\underline{190}$ $\underline{22}$. $\underline{15}$ $\underline{27}$ $\underline{18}$ $\underline{5}$ $\underline{15}$ |
|---|---|
| Site Sketch Woone | rockitcisper) KN |
| FLOW | CHEEK CHEEK |
| 1+ 0300W | |



Notes (Photo Number and substrate description):

Coold's collected in contex of transect

Desand, look/grance

Desand + fine grand

Stand + fine grand

Stand + fine grand

67 - looky PS

Grand (arse) + could

Stand


Stand

Bedroch

Stand

Bedroch

Sediment Probing Section Form




Notes (Photo Number and substrate description):

Coords collected from center of transect While board

- O hed woon

- (5) fine silt on hed wich A Co and in bud will
- DSut m/ smart on bedwell SSIIt on bedwell
- 6 silty clay
- 10 Elay on molicol




| | Sample Location | | | | | | | rs:5W3 | | - 08(8 |
|---------|--|-------------|----------------|--------------|--------------|---------|-----------|----------------------------|------------------------------------|------------------|
| 30 - | Bank Width (m)_ Sediment Depths | | measu 17.5" | ATTACK AND | — Num 35' | 45' | diment De | epth Estim 58' | ates: [| 1 72' 8.5' |
| WD: | 4.5" 5.0" 12.5" Site Sketch | 13.011 | 8" | - 1811 F" | - Du | 25' | 7" | 15" | 15" | 20,5" |
| SD=redu | net | | | TV | celine | | 2 FALLS | KD_ | K | N |
| WD = wa | | ion | | | | | | AT RIVE CREEK Thar w |) | |
| | | + 4 | V | CITY | OF PAR | x x x | x x | × U × | -fen | celile |
| | Cross Section (p | | | izontal ar | nd vertical | measure | ments) | . (| R.B | |
| | | | \$ | | 6 3 | | | | -mudi Sedim depth (refusa | ent |
| | Notes (Photo Nu | | | | | u al | tra | | | |
| | Coords 51 - White bo 52 - Tooking 53 - Tooking D Silty Sand D Sand + Silt B) Glandy 9m | DS W Som | 54 55 56 | -looky | D proba | ng | , , , , , | | | |



| | Sample Location | | | | | | | | | |
|--------------------|-----------------------------|---|---|---------------------|------------|--------------|------------|-----------------------|---------|-----------|
| | Longitude: 3 | 1.13.160 | 1606 10 | | atitude | e: <u>90</u> | ,54023 | 7/21°W | | |
| Codes | Bank Width (m)_ | 113 | ft. | N N | umber | of Sedi | ment De | oth Estima | ates:10 |) |
| cen | Sediment Depth | s: (5 to 10 i | measuremen | nts) 56 | | Less remend | 781 | 991 | - | |
| SSP | 57.5 5 7.5" | 3.0" | >60+" > | 260" 47 4.0" 15. | 04 | 260" | 35" | 29" | 0 3 | , bedrook |
| | Site Sketch | 3.0 | | Cliff | | art | 33 . | 43. | ,,,, | |
| nd dopte | | 8 | ing River | Flow | | - | | | | |
| Japth | | | 2 | | | 1 | | | | |
| | | | | | | | | | | |
| | | | | | 1 | | | | | |
| | | 5.1 | Bar | | | 1 | | | | |
| | | SUND | 041 | | | 1 | - | | | |
| | | | | | | 1 | | | | |
| | 7 | Back | Water | | | | | | | |
| | | | | 4 | | 1 | | | | |
| | | | and the second second second | | Polumenton | | | | | |
| | Mag Not 2 Cross Section (p | | | al and verti | | easureme | ents) | | | |
| ted K | | | | | | easureme | ents) | 10 | | Bedra |
| ted K RB | | | | | | easureme | ents) | 10 | | Bedra Chi |
| ted K RB | Cross Section (p | mber and | h horizonta | al and verti | cal me | | 4 | | | L B |
| ted RB | Cross Section (p | rovide bot | substrate d | al and verti | cal me | | a lever | ing D.S. | | L B |
| ted K RB | Notes (Photo Nu | mber and white be | substrate de trans. | escription) | cal me | | a lever | | | L B |
| ted K RB | Notes (Photo Nu | mber and white be D.S. at D.S. at Loweing | substrate d | escription) | cal me | | a lever | ing D.S. | | L B |
| Da | Notes (Photo Nu | mber and white be as a boston, proper | substrate de trans. I trans. Sedinet fo | escription) | cal me | 117- | Levo R | ing D.S. | | L B |
| Da | Notes (Photo Nu | mber and white be as a boston, proper | substrate de trans. I trans. Sedinet fo | escription) | cal me | | Levo R | ing D.S. | | L B |
| Do (1) f | Notes (Photo Nu | mber and White be D.S. ed Louking D.S. on Fronze | substrate de sard trans. I trans. Sedinet from Sedinet | escription) | cal me | 117- | Levo R | ing D.S. | | L B |
| Da 1) 6 2) | Notes (Photo Nu | mber and white be D.S. at D.S. at D.S. probe | substrate de sard trans. I trans. Sedinet from Sedinet | escription) | cal me | 117- | Levo R | ing D.S. | | L B |
| Da (1) (6 2) 3 (4) | Notes (Photo Nu | mber and White be D.S. at D.S. at Looking D.S. on for | substrate de sard trans. I trans. Sedinet from Sedinet | escription) | cal me | 117- | Leve above | ing D.S. Fransect. | | L B |





bed rock



| | | | Date: 121112 Samplers: 35, 50, 81 | |
|-------------|------------------------|----------------------------|---|---------------------------|
| | Longitude: | 37.95432255°N | Latitude: 90,53966347°W | |
| D Inciement | Bank Width (m) | 120.5 FL | Number of Sediment Depth Estimates: 10 | Every 12 f |
| | Sediment Depths | : (5 to 10 measurements) | 66 | . From RO to |
| nch 60. | 24" 8.5" | 4.0" 6.0" 23" | 41" 16" 11" 29" 38" | |
| (2n) 54 50 | 0" 0" | .0" 0" 0" | 41" 16" 11" 29" 38" 5.5" LB 12.0" 23.0" 29.5" 27.5" | |
| NO | Site Sketch | Cliff | | |
| | | | | |
| | | Bis River | \rightarrow | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | Gravel / Sand Bar | | () |
| | | t the second | | (wester Rout Wed |
| | | | | Root Wed tree (LWO) |
| | | | | (2000) |
| | | | | |
| | | RB | | |
| | | | | |
| | Cross Section (p | rovide both horizontal and | vertical measurements) | |
| | 1 2 | 13 4 | 15 16 17-10 | |
| | | | 19 | 10 |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | mber and substrate descrip | tion): | |
| | 125 ~ white | | | |
| | 126 - U.S. 127 D.S. | @ transect | | |
| | 128 LSen | | t transect | |
| | 124 Lak | - | transect | |
| | | | | |
| | 1) Fire grand I san |) | same as about | |
| | 21 | 7) | Fine Scorel (Sen) (Silt | |
| | 3) -, ~ | 8) | | |
| | 4) ~ " | 9 | | |
| | | | | |



